

**Daily interventions (continued)**

Please record all interventions given on each day of admission using a cross, unless otherwise specified.

An item should be recorded in the PCCMDS when the critical care activity applies for a period of greater than 4 hours.

If no interventions given select 'No defined critical care activity'

Admission date: .....



|  |   | Day     | 0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 |
|--|---|---------|---|---|---|---|---|---|---|---|---|---|----|----|----|----|
| <b>Basic</b>   | No defined critical care activity   | Code 99 |   |   |   |   |   |   |   |   |   |   |    |    |    |    |
|  | Continuous ECG monitoring   | 50      |   |   |   |   |   |   |   |   |   |   |    |    |    |    |
|  | Continuous pulse oximetry   | 73      |   |   |   |   |   |   |   |   |   |   |    |    |    |    |
| <b>Airway and ventilatory</b>  | Invasive ventilation via endotracheal tube  | 51      |   |   |   |   |   |   |   |   |   |   |    |    |    |    |
|  | Invasive ventilation via tracheostomy tube  | 52      |   |   |   |   |   |   |   |   |   |   |    |    |    |    |
|  | Non-invasive ventilatory support  | 53      |   |   |   |   |   |   |   |   |   |   |    |    |    |    |
|  | Advanced ventilatory support (jet ventilation)  | 56      |   |   |   |   |   |   |   |   |   |   |    |    |    |    |
|  | Advanced ventilatory support (oscillatory ventilation)                                  | 56      |   |   |   |   |   |   |   |   |   |   |    |    |    |    |
|  | Nasopharyngeal airway   | 55      |   |   |   |   |   |   |   |   |   |   |    |    |    |    |
|  | Tracheostomy cared for by nursing staff   | 13      |   |   |   |   |   |   |   |   |   |   |    |    |    |    |
|  | Supplemental oxygen therapy (irrespective of ventilatory state)                         | 09      |   |   |   |   |   |   |   |   |   |   |    |    |    |    |
|  | Maximal oxygen concentration (record maximum concentration as %)                        | -       |   |   |   |   |   |   |   |   |   |   |    |    |    |    |
|  | High flow nasal cannula therapy (record maximum daily flow in L/minute)                 | 88      |   |   |   |   |   |   |   |   |   |   |    |    |    |    |
|  | Upper airway obstruction requiring nebulised adrenaline (epinephrine)                   | 57      |   |   |   |   |   |   |   |   |   |   |    |    |    |    |
|  | Apnoea requiring intervention (>3 in 24 hours or need for bag-mask ventilation)         | 58      |   |   |   |   |   |   |   |   |   |   |    |    |    |    |
|  | Acute severe asthma requiring IV bronchodilator therapy or continuous nebuliser         | 59      |   |   |   |   |   |   |   |   |   |   |    |    |    |    |
| Unplanned extubation (record number of unplanned extubations)                | 90  |         |   |   |   |   |   |   |   |   |   |   |    |    |    |    |
| Unplanned tracheostomy removal or change (record number of unplanned events) | -   |         |   |   |   |   |   |   |   |   |   |   |    |    |    |    |
| <b>Cardio-vascular</b>   | Arterial line monitoring  | 60      |   |   |   |   |   |   |   |   |   |   |    |    |    |    |
|  | External pacing   | 61      |   |   |   |   |   |   |   |   |   |   |    |    |    |    |
|  | Central venous pressure monitoring  | 62      |   |   |   |   |   |   |   |   |   |   |    |    |    |    |
|  | Continuous infusion of inotrope, vasodilator or prostaglandin                           | 06      |   |   |   |   |   |   |   |   |   |   |    |    |    |    |
|  | Bolus IV fluids (>80 ml/kg/day) in addition to maintenance IV fluids                    | 63      |   |   |   |   |   |   |   |   |   |   |    |    |    |    |
|  | Cardio-pulmonary resuscitation  | 64      |   |   |   |   |   |   |   |   |   |   |    |    |    |    |
|  | Extracorporeal membrane oxygenation (ECMO)  | 65      |   |   |   |   |   |   |   |   |   |   |    |    |    |    |
|  | Ventricular assist device (VAD)   | 65      |   |   |   |   |   |   |   |   |   |   |    |    |    |    |
|  | Aortic balloon pump   | 65      |   |   |   |   |   |   |   |   |   |   |    |    |    |    |
| Arrhythmia requiring intravenous anti-arrhythmic therapy                     | 94  |         |   |   |   |   |   |   |   |   |   |   |    |    |    |    |
| <b>Renal</b>   | Peritoneal dialysis   | 05      |   |   |   |   |   |   |   |   |   |   |    |    |    |    |
|  | Haemofiltration   | 16      |   |   |   |   |   |   |   |   |   |   |    |    |    |    |
|  | Haemodialysis   | 66      |   |   |   |   |   |   |   |   |   |   |    |    |    |    |
|  | Plasma filtration   | 67      |   |   |   |   |   |   |   |   |   |   |    |    |    |    |
|  | Plasma exchange   | 67      |   |   |   |   |   |   |   |   |   |   |    |    |    |    |
| <b>Neuro-logical</b>   | ICP-intracranial pressure monitoring  | 68      |   |   |   |   |   |   |   |   |   |   |    |    |    |    |
|  | Intraventricular catheter or external ventricular drain                                 | 69      |   |   |   |   |   |   |   |   |   |   |    |    |    |    |
|  | Status epilepticus requiring treatment with continuous infusion of anti-epileptic drugs | 97      |   |   |   |   |   |   |   |   |   |   |    |    |    |    |
|  | Reduced conscious level (GCS ≤ 12) AND hourly (or more frequent) GCS monitoring         | 95      |   |   |   |   |   |   |   |   |   |   |    |    |    |    |
| <b>Analgesia/sedation</b>  | Epidural catheter in situ   | 85      |   |   |   |   |   |   |   |   |   |   |    |    |    |    |
|  | Continuous intravenous infusion of a sedative agent                                     | 96      |   |   |   |   |   |   |   |   |   |   |    |    |    |    |
| <b>Metabolic</b>   | Diabetic ketoacidosis (DKA) requiring continuous infusion of insulin                    | 70      |   |   |   |   |   |   |   |   |   |   |    |    |    |    |
| <b>Other</b>   | Exchange transfusion  | 04      |   |   |   |   |   |   |   |   |   |   |    |    |    |    |
|  | Intravenous thrombolysis  | 71      |   |   |   |   |   |   |   |   |   |   |    |    |    |    |
|  | Extracorporeal liver support using molecular absorbent recirculating system (MARS)      | 72      |   |   |   |   |   |   |   |   |   |   |    |    |    |    |
|  | Patient nursed in single occupancy cubicle (state reason for isolation below)           | 74      |   |   |   |   |   |   |   |   |   |   |    |    |    |    |
| <b>High cost drugs</b>   | Medical gases Band 1 – nitric oxide   | X841    |   |   |   |   |   |   |   |   |   |   |    |    |    |    |
|  | Surfactant  | X842    |   |   |   |   |   |   |   |   |   |   |    |    |    |    |

Reason for isolation (if patient nursed in single occupancy cubicle)

Contact us · picanet@leeds.ac.uk · 0113 343 8125

For more contact details, go to [www.picanet.org.uk/contact-us](http://www.picanet.org.uk/contact-us)

For forms, dataset manuals and guidance, go to <https://www.picanet.org.uk/level-2-expansion/>

